

Title: **Retinoscopy/Autorefraction, which is the best starting point for a non-cycloplegic refraction?**

Running Title: Retinoscopy vs. Autorefraction

Authors: J Jorge¹, MSc, Member of faculty

 A Queirós¹, Member of faculty

 JB Almeida¹, MSc, PhD, Member of faculty

 MA Paraíta², MSc, MD, PhD, Member of faculty

Institutions: 1 Department of Physics (Optometry), School of Sciences. University of Minho. Braga. Portugal.

 2 Department of Surgery (Ophthalmology), School of Optics and Optometry. University of Santiago de Compostela. Spain

Corresponding Author: Jorge Jorge

Address: Departamento de Física,
 Universidade do Minho
 Campus de Gualtar
 4710 – 057 Braga
 Portugal

Tel: +351 253 604 333

Fax: +351 253 604 061

E-mail: jorge@fisica.uminho.pt

The authors state that they have no proprietary or commercial interest in Autorefractor Nidek ARK 700A.

Key words: Refraction, refractive error, accuracy, automated refractor, retinoscopy, subjective refraction, orthogonal functions, and astigmatism.

Acknowledgment: Authors thank contributions of José Manuel González-Méijome.

ABSTRACT

Purpose: The aim of this study was to estimate the agreement between an autorefractor (Nidek ARK 700A) and retinoscopy with subjective refraction.

Methods: Measurements of autorefraction obtained with the ARK700A and retinoscopy were performed on 192 right eyes from 192 healthy young adults and compared with subjective refraction. These measurements were performed without cycloplegia. The age range was 18-34 years, with a mean value of 21.6 years and a standard deviation (SD) of 2.66 years.

Results: A comparison of the autorefractor and subjective refraction results shows that; (1) for the mean spherical equivalent (M) the autorefractor yields more negative values (-0.44 ± 0.54 D, $p = 0.000$); (2) for the Jackson cross-cylinder at axis 0° (J_0), the autorefractor yields more positive values than the subjective ones (-0.05 ± 0.13 D, $p = 0.000$); (3) and for the Jackson cross-cylinder at axis 45° (J_{45}), the autorefractor results are more negative (-0.02 ± 0.09 D, $p = 0.019$). The differences found for each component M, J_0 e J_{45} are statistically significant.

By comparing retinoscopy with the subjective exam, there are no statistically significant differences found for the M component, (-0.02 ± 0.33 D, $p = 0.304$). For the J_0 and J_{45} components, the differences are statistically significant (-0.07 ± 0.10 D, $p = 0.000$; -0.01 ± 0.08 D, $p = 0.008$).

Conclusions: the present results confirm that when performed by an experienced clinician, retinoscopy is more accurate than automatic refraction giving a better starting point to non-cycloplegic refraction.